

University of Groningen

Practice patterns in osteoporosis prevention in patients on glucocorticoids

Naunton, M; Duyvendak, M; Peterson, GM; Brouwers, JRBJ

Published in:
Osteoporosis International

DOI:
[10.1007/s00198-005-0055-0](https://doi.org/10.1007/s00198-005-0055-0)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Naunton, M., Duyvendak, M., Peterson, GM., & Brouwers, JRBJ. (2006). Practice patterns in osteoporosis prevention in patients on glucocorticoids. *Osteoporosis International*, 17(4), 634-635.
<https://doi.org/10.1007/s00198-005-0055-0>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Practice patterns in osteoporosis prevention in patients on glucocorticoids

M. Naunton · M. Duyvendak · G. M. Peterson ·
J. R. B. J. Brouwers

Published online: 1 March 2006

© International Osteoporosis Foundation and National Osteoporosis Foundation 2006

Dear Editor,

We have read the recent results from the study on osteoporosis prevention in chronic (>90 days) glucocorticoid users by Feldstein et al. [1] and have three comments.

1. We believe that any apparent under-treatment may be partly due to the fact that the data analysed were collected from January 2000–December 2001. The American College of Rheumatology (ACR) guidelines for prevention of glucocorticoid-induced osteoporosis were published in July 2001 and recommended that osteoporosis prevention should be considered in those patients treated on ≥ 5 mg of

prednisolone per day [3]. The guidelines available before this suggested that patients should be considered for osteoporosis prevention if the dose was greater than 7.5 mg per day, and the patients were treated for >6 months with the glucocorticoid [4]. We believe to be fair to the prescribers and other primary-care clinicians it should be noted that at the time that these data were collected most would only be aware that patients treated with ≥ 7.5 mg prednisolone should be considered for osteoporosis prevention.

2. There was no information on calcium or vitamin D (activated or non-activated) use. Were these medications/supplements excluded/not considered or unavailable? We believe these are important to mention because they were listed in the 1996 and 2001 ACR guidelines. In addition, clinical trials with bisphosphonates have included calcium/vitamin D in their protocol.

3. The authors note that their population had a high mean prednisone dose of 20mg/day. We believe that this unusually high dose of corticosteroid is due partly to their methods in calculating the “average” dose. The mean dose was obtained by using supplied quantities and not the actual dose. The authors included all medications dispensed including those during hospitalisation, which probably accounts for the high mean dose. For example, asthmatic/COPD patients (or other patients with sliding scales) with frequent admissions to hospital are likely to receive short courses of high-dose corticosteroid therapy in hospital, which will undoubtedly increase the mean dose. It would be interesting to know if these doses were excluded, and what the median (and range) dose of corticosteroid therapy was, i.e. what is the usual maintenance dose of corticosteroid? This may be an important consideration for prescribers when considering osteoporosis prevention because the correlation between daily dose and fracture is stronger than for cumulative dose [2]. Having said that, we don’t expect the overall results from this study (i.e., underuse of osteoporosis prevention) would change significantly.

M. Naunton (✉)
Department of Pharmacotherapy and Pharmaceutical Care,
State University of Groningen,
Antonius Deusinglaan 1,
9713 AV Groningen, The Netherlands
e-mail: M.Naunton@rug.nl
Tel.: +31-503-637576
Fax: +31-503-632772

M. Duyvendak
Department of Clinical Pharmacy, Tjongerschans Hospital,
Postbus 10500, 8440 MA Heerenveen, The Netherlands
e-mail: mduyvendak@tjongerschans.nl
Tel.: +31-513-685693
Fax: +31-513-685945

G. M. Peterson
Unit for Medication Outcomes, Research and Education,
Tasmanian School of Pharmacy,
Hobart, 7000, Tasmania, Australia
e-mail: G.Peterson@utas.edu.au
Tel.: +61-62-262203

J. R. B. J. Brouwers
Department of Pharmacotherapy and Pharmaceutical Care,
State University of Groningen,
Antonius Deusinglaan 1,
9713 AV Groningen, The Netherlands
e-mail: J.R.B.J.Brouwers@rug.nl
Tel.: +31-503-637576

References

1. Feldstein AC, Elmer PJ, Nichols GA, Herson M (2005) Practice patterns in patients at risk for glucocorticoid-induced osteoporosis. *Osteoporos Int* Sep 3 [Epub ahead of print]
2. van Staa TP, Leufkens HG, Abenhaim L, Zhang B, Cooper C (2000) Oral corticosteroids and fracture risk: relationship to daily and cumulative doses. *Rheumatology* 39:1383–1389
3. Recommendations for the prevention and treatment of glucocorticoid-induced osteoporosis: 2001 update (2001) American college of rheumatology ad hoc committee on glucocorticoid-induced osteoporosis. *Arthritis Rheum* 44(7):1496–1503
4. Recommendations for the prevention and treatment of glucocorticoid-induced osteoporosis (1996) American college of rheumatology task force on osteoporosis guidelines. *Arthritis Rheum* Nov 39(11):1791–1801